



POLICY: Oncology (Injectable – Bispecific – CD20-Directed) – Epkinly Utilization Management Medical Policy

- Epkinly™ (epcoritamab-bysp subcutaneous injection – Genmab)

EFFECTIVE DATE: 11/15/2023

LAST REVISION DATE: 06/04/2025; selected revisions 12/03/2025 and 01/28/2026

COVERAGE CRITERIA FOR: All UCare Plans

OVERVIEW

Epkinly, a bispecific CD20-directed CD3 T-cell engager, is indicated for the treatment of:

- Relapsed or refractory **diffuse large B-cell lymphoma (DLBCL)**, not otherwise specified, including DLBCL arising from indolent lymphoma, and high-grade B-cell lymphoma, in adults after two or more lines of systemic therapy.¹
- Relapsed or refractory follicular lymphoma (FL) in adults after two or more lines of systemic therapy.¹
- Relapsed or refractory follicular lymphoma (FL) in adults, in combination with lenalidomide and rituximab.¹

Dosing Information

Epkinly is administered by subcutaneous injection.¹ Table 1, 2 and 3 summarize the dosing schedule for Epkinly.

Table 1. Epkinly Dosing Schedule for DLBCL and High-grade B-cell Lymphoma.¹

Treatment Cycle	Treatment Day	Dose
Cycle 1	Day 1	0.16 mg
	Day 8	0.8 mg
	Day 15	48 mg
	Day 22	48 mg
Cycle 2 and 3	Days 1, 8, 15, 22	48 mg
Cycle 4 to 9	Days 1 and 15	48 mg
Cycle 10+	Day 1	48 mg

Table 2. Epkinly Dosing Schedule for Follicular Lymphoma¹

Treatment Cycle	Treatment Day	Dose
Cycle 1	Day 1	0.16 mg
	Day 8	0.8 mg
	Day 15	3 mg
	Day 22	48 mg
Cycle 2 and 3	Days 1, 8, 15, 22	48 mg
Cycle 4 to 9	Days 1 and 15	48 mg
Cycle 10+	Day 1	48 mg

Table 3. Epkinly Dosing Schedule in Combination with lenalidomide and rituximab for Follicular Lymphoma¹

Treatment Cycle	Treatment Day	Dose
Cycle 1	Day 1	0.16 mg
	Day 8	0.8 mg
	Day 15	3 mg

	Day 22	48 mg
Cycles 2 and 3	Days 1, 8, 15, 22	48 mg
Cycles 4 to 12	Day 1	48 mg

Guidelines

Epkinly has been addressed by National Comprehensive Cancer Network (NCCN) guidelines:

- **B-cell lymphoma** clinical practice guidelines (version 1.2026 – December 22, 2025) recommend Epkinly for second-line and subsequent treatment of classic follicular lymphoma, DLBCL, high-grade B-cell lymphomas, human immunodeficiency virus (HIV)-related B-cell lymphomas, and post-transplant lymphoproliferative disorders. Epkinly is also recommended as single agent use for histologic transformation of indolent lymphomas to DLBCL after at least two lines of systemic therapy.^{2,3}
- **Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma** clinical practice guidelines (version 2.2026 – December 22, 2025) recommend Epkinly for the treatment of histologic Richter transformation (category 2A).⁴

Safety

Epkinly has Boxed Warnings for cytokine release syndrome and immune effector cell-associated neurotoxicity syndrome.¹

POLICY STATEMENT

Prior Authorization is recommended for medical benefit coverage of Epkinly. Approval is recommended for those who meet the **Criteria** and **Dosing** for the listed indication. Extended approvals are allowed if the patient continues to meet the Criteria and Dosing. Requests for doses outside of the established dosing documented in this policy will be considered on a case-by-case basis by a clinician (i.e., Medical Director or Pharmacist). All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with Epkinly as well as the monitoring required for adverse events and long-term efficacy, approval requires Epkinly to be prescribed by or in consultation with a physician who specializes in the condition being treated.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Epkinly is recommended in those who meet the following criteria:

FDA-Approved Indication

1. **Diffuse Large B-Cell Lymphoma.** Approve for 1 year if the patient meets ALL of the following (A, B, and C):

Note: Diffuse large B-cell lymphoma (DLBCL) includes DLBCL not otherwise specified, DLBCL arising from indolent lymphoma, and high-grade B-cell lymphoma.

A) Patient is ≥ 18 years of age; AND

B) Patient has received one or more lines of systemic therapy; AND

Note: Examples of systemic therapy include RCHOP (rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone) and DHA (dexamethasone, cytarabine) + platinum (carboplatin, cisplatin, or oxaliplatin) \pm rituximab.

C) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve the following dosing regimen (A and B):

- A) The dose is up to 48 mg administered by subcutaneous injection; AND
- B) The agent is given in 28-day cycles that meet the following (i, ii, and iii):
 - i. Cycles 1, 2, and 3: Maximum of 4 injections; AND
 - ii. Cycles 4 to 9: Maximum of 2 injections; AND
 - iii. Cycles 10 and beyond: maximum of 1 injection.

2. **Classic Follicular Lymphoma.** Approve for 1 year if the patient meets ALL of the following (A, B, and C):

- A) Patient is ≥ 18 years of age; AND
- B) Patient has received one or more lines of systemic therapy; AND
Note: Examples of systemic therapy include CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) plus rituximab or Gazyva (Obinutuzumab intravenous infusion) and CVP (cyclophosphamide, vincristine, prednisone) plus rituximab or Gazyva.
- C) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve the following dosing regimen (A and B):

- A) The dose is up to 48 mg administered by subcutaneous injection; AND
- B) The agent is given in 28-day cycles that meet the following (i, ii, and iii):
 - i. Cycles 1, 2, and 3: Maximum of 4 injections; AND
 - ii. Cycles 4 to 9: Maximum of 2 injections; AND
 - iii. Cycles 10 and beyond: maximum of 1 injection.

Other Uses with Supportive Evidence

3. **Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma.** Approve for 1 year if the patient meets ALL of the following (A, B, and C):

- A) Patient is ≥ 18 years of age; AND
- B) Patient has histologic Richter transformation; AND
- C) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve the following dosing regimen (A and B):

- A) The dose is up to 48 mg administered by subcutaneous injection; AND
- B) The agent is given in 28-day cycles that meet the following (i, ii, and iii):
 - i. Cycles 1, 2, and 3: Maximum of 4 injections; AND
 - ii. Cycles 4 to 9: Maximum of 2 injections; AND
 - iii. Cycles 10 and beyond: maximum of 1 injection

4. **Human Immunodeficiency Virus (HIV)-Related B-Cell Lymphomas.** Approve for 1 year if the patient meets ALL of the following (A, B, and C):

Note: HIV-related B-cell lymphomas includes HIV-related diffuse large B-cell lymphoma (DLBCL), primary effusion lymphoma, and human herpes virus-8 (HHV8) positive DLBCL.

- A) Patient is ≥ 18 years of age; AND
 - B) Patient has received one or more lines of systemic therapy; AND
Note: Examples of systemic therapy include RCHOP (rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone) and R-EPOCH (rituximab, etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin).
 - C) The medication is prescribed by or in consultation with an oncologist.
-

Dosing. Approve the following dosing regimen (A and B):

- A) The dose is up to 48 mg administered by subcutaneous injection; AND
- B) The agent is given in 28-day cycles that meet the following (i, ii, and iii):
 - i. Cycles 1, 2, and 3: Maximum of 4 injections; AND
 - ii. Cycles 4 to 9: Maximum of 2 injections; AND
 - iii. Cycles 10 and beyond: maximum of 1 injection.

5. Post-Transplant Lymphoproliferative Disorders. Approve for 1 year if the patient meets ALL of the following (A, B, and C):

- A) Patient is ≥ 18 years of age; AND
- B) Patient has received one or more lines of systemic therapy; AND
Note: Examples of systemic therapy include RCHOP (rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone) and RCEPP (rituximab, cyclophosphamide, etoposide, prednisone, procarbazine).
- C) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve the following dosing regimen (A and B):

- A) The dose is up to 48 mg administered by subcutaneous injection; AND
- B) The agent is given in 28-day cycles that meet the following (i, ii, and iii):
 - i. Cycles 1, 2, and 3: Maximum of 4 injections; AND
 - ii. Cycles 4 to 9: Maximum of 2 injections; AND
 - iii. Cycles 10 and beyond: maximum of 1 injection.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Epkinly is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Epkinly subcutaneous injection [prescribing information]. Plainsboro, NJ: Genmab; November 2025.
2. The NCCN Drugs and Biologics Compendium. © 2025 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on January 22, 2026. Search term: eporitamab.
3. The NCCN B-Cell Lymphomas Clinical Practice Guidelines in Oncology (version 1.2026 – December 22, 2025). © 2025 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on January 22, 2026.
4. The NCCN Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma Clinical Practice Guidelines in Oncology (version 2.2026 – December 22, 2025). © 2025 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on January 22, 2026.

HISTORY

Type of Revision	Summary of Changes	Review Date
New Policy	--	06/05/2023
Annual Revision	Classic Follicular Lymphoma: Added new condition of approval.	06/12/2024
UCare P&T Review	Policy reviewed and approved by UCare P&T committee. Annual review process	09/16/2024
Annual Revision	Diffuse Large B-Cell Lymphoma, Human Immunodeficiency Virus (HIV)-Related B-Cell Lymphomas, and Post-Transplant Lymphoproliferative Disorders :	06/04/2025

	Removed the requirement that the medication is used as a single agent. For the requirement that patient has received two or more lines of systemic therapy, this was changed to one or more lines of systemic therapy. Classic Follicular Lymphoma: Moved Classic Follicular Lymphoma from “Other Uses with Supportive Evidence” to “FDA-Approved Indication”.	
UCare P&T Review	Policy reviewed and approved by UCare P&T committee. Annual review process	09/15/2025
Selected Revision	Classic Follicular Lymphoma: The requirement that patient has received two or more lines of systemic therapy was modified to patient has received one or more lines of systemic therapy. The requirement that the medication is given as a single agent was removed.	12/03/2025
Selected Revision	Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma: Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma was added as a new condition for approval.	01/28/2026