

POLICY: Dermatology – Ycanth Utilization Management Medical Policy

- Ycanth™ (cantharidin 0.7% topical solution – Verrica)

EFFECTIVE DATE: 07/01/2024**REVIEW DATE:** 02/19/2025; selected revision 03/12/2025**COVERAGE CRITERIA FOR:** ALL UCARE PLANS

OVERVIEW

Ycanth, a cantharidin-based topical solution, is indicated for the treatment of molluscum contagiosum in patients 2 years of age and older.¹

Disease Overview

Molluscum contagiosum is a viral skin infection of the Poxviridae family that can cause white, pink, or flesh colored bumps either alone, or in groups; it is spread by direct contact.^{2,3} Common locations are the trunk, face, and extremities. Patients may experience pain, itching, and eczema, as well as secondary bacterial infections. Resolution usually occurs within 6 to 12 months; in selected cases it can take longer for the skin infection to completely disappear. The condition is found in children and adults; however, it is more common in younger patients. Immunocompetent patients can often clear the infection without treatment. However, patients with additional dermatologic conditions (e.g., atopic dermatitis), or in those who are immunocompromised, have more extensive infection that is harder to treat. Molluscum contagiosum is most common in warm, humid climates.

Clinical Efficacy

The efficacy of Ycanth for the treatment of molluscum contagiosum infections has been evaluated in two pivotal studies.^{1,4} The studies included patients ≥ 2 years of age with a clinical diagnosis of molluscum contagiosum with treatable lesions. The primary efficacy endpoint was the proportion of the Ycanth treated patients achieving complete clearance of all molluscum contagiosum lesions compared to those who received the vehicle at Day 84 of trial.

Guidelines

The American Academy of Pediatrics (AAP) RedBook 2024-2027 cite that cryotherapy, curettage, and cantharidin (compounded or Ycanth) have the most support for treatment.²

POLICY STATEMENT

Prior Authorization is recommended for medical benefit coverage of Ycanth. Approval is recommended for those who meet the **Criteria** and **Dosing** for the listed indication. Extended approvals are allowed if the patient continues to meet the Criteria and Dosing. Requests for doses outside of the established dosing documented in this policy will be considered on a case-by-case basis by a clinician (i.e., Medical Director or Pharmacist). All approvals are provided for the

duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Ycanth is recommended in those who meet the following criteria:

FDA-Approved Indication

1. **Molluscum Contagiosum.** Approve Ycanth for 3 months if the patient meets ALL of the following (A, B, and C):
 - A) Patient is ≥ 2 years of age; AND
 - B) Approve if patient meets ONE of the following (i or ii):
 - i. Patient is treating new lesions that have not previously been treated with Ycanth; OR
 - ii. Patient is treating lesions that have been previously treated with Ycanth for less than 4 treatment cycles; AND
 - C) Ycanth is being administered by a healthcare professional.

Dosing. Approve up to two applicators per treatment, not more frequently than once every 21 days.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Ycanth is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Ycanth™ topical solution [prescribing information]. West Chester, PA: Verrica; October 2024.
2. American Academy of Pediatrics. Red Book: 2024-2027 report of the Committee of Infectious Diseases (33rd Edition). Molluscum Contagiosum. Pages 603-604.
3. Badri T, Gandhi GR. Molluscum Contagiosum. [Updated March 27, 2023]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK441898/>. Accessed on February 17, 2025.
4. Eichenfield LF, McFalda W, Barbec B, et al. Safety and Efficacy of VP-102, a proprietary, drug-device combination product containing cantharidin, 0.7% (w/v), in children and adults with molluscum contagiosum: two phase 3 randomized clinical trials. *JAMA Dermatol.* 2020;156(12):1315-1323.

HISTORY

Type of Revision	Summary of Changes	Review Date
New Policy		02/28/2024
UCare P&T Review	Policy reviewed and approved by UCare P&T committee. Annual review process	09/16/2024
Annual Revision	No criteria changes.	02/19/2025

Selected Revision	Molluscum Contagiosum: Dosing was updated to approve up to two applicators per treatment, not more frequently than once every 21 days.	03/12/2025
----------------------	---	------------