

POLICY: Hemophilia – Gene Therapy – Beqvez Utilization Management Medical Policy

- Beqvez™ (fidanacogene elaparovvec-dzkt intravenous infusion – Pfizer)

EFFECTIVE DATE: 8/15/2024

LAST REVISION DATE: 02/26/2025

COVERAGE CRITERIA FOR: ALL UCARE PLANS

OVERVIEW

Beqvez, an adeno-associated virus (AAV) vector-based gene therapy, is indicated for the treatment of **hemophilia B** (congenital Factor IX deficiency) in adults with moderate to severe disease who: 1) currently use Factor IX prophylaxis therapy; or 2) have current or historical life-threatening hemorrhage; or 3) have repeated, serious spontaneous bleeding episodes, AND do not have neutralizing antibodies to adeno-associated virus serotype Rh74var (AAVRh74var) capsid as detected by an FDA-approved test.^{1,2} The recommended dose of Beqvez is 5×10^{11} vector genomes per kg of body weight given as a one-time (per lifetime) single dose as an intravenous infusion.¹ Dose based on adjusted body weight for those with a body mass index $> 30 \text{ kg/m}^2$.

Disease Overview

Hemophilia B is a genetic bleeding disorder caused by missing or insufficient levels of blood Factor IX, a protein required to produce blood clots to halt bleeding.³⁻⁶ The condition is a rare X-linked bleeding disorder that mainly impacts males. Hemophilia B is four times less common than hemophilia A, which is caused by a relative lack of blood Factor VIII. Approximately 30,000 individuals are living with hemophilia in the US and hemophilia B accounts for around 15% to 20% of hemophilia cases, or around 6,000 patients. Symptoms include heavy or prolonged bleeding following an injury or after a medical procedure. Bleeding can also occur internally into joints, muscles, or internal organs. Spontaneous bleeding events may also occur. Complications in patients with hemophilia B include joint disease and hemarthrosis. Hemophilia B may be diagnosed when bleeding occurs in infancy or later in life for those with milder disease. There is a strong correlation between Factor IX levels and phenotypic expression of bleeding. Normal plasma levels of Factor IX range from 50% to 150%. The disease is classified based on reduced levels. Mild, moderate, and severe hemophilia B is characterized by Factor IX levels ranging from 6% up to 49%, 1% up to 5%, and $< 1\%$, respectively. Besides gene therapies for the treatment of hemophilia B, Factor IX products, both recombinant and plasma-derived, are used routinely to prevent bleeding or are given on-demand to treat bleeding episodes associated with hemophilia B.³⁻⁶ Hympavzi™ (marstacimab subcutaneous injection), a tissue factor pathway inhibitor antagonist, is also FDA approved for routine prophylaxis to prevent or reduce the frequency of bleeding episodes in adult and pediatric patients ≥ 12 years of age with hemophilia B (congenital Factor IX deficiency) without inhibitors; it is also indicated for use in hemophilia A.⁷

Clinical Efficacy

The efficacy of Beqvez was evaluated in one ongoing, prospective, open-label, single-arm, single-dose, multinational, Phase III pivotal trial called BENEENE-2 involving adult males with moderately severe to severe hemophilia B (Factor IX activity $\leq 2\%$) [$n = 45$].^{1,2} All patients completed a prospective lead-in period of at least 6 months in which baseline data were collected while patients were receiving Factor IX products for routine prophylaxis.¹ However, after receipt of Beqvez, use of such products for routine prophylaxis was to be suspended. The trial is ongoing with a planned long-term follow-up of 6 years. Patients were required to be negative for pre-existing neutralizing antibodies to AAVRh74var capsid to

participate. Factor IX inhibitors (or a history), uncontrolled human immunodeficiency virus (HIV) infection, or significant liver fibrosis were exclusion criteria. Adequate hepatic and renal function were required. The median follow-up was 2.0 years (range 0.4 to 3.2 years) post-Beqvez administration. The model-derived mean annualized bleeding rate was 4.5 bleeds/year during the baseline lead-in period vs. 2.5 bleeds/year during the post-Beqvez efficacy evaluation period. In total, 60% of patients did not experience any bleeds after receipt of Beqvez; only 29% of patients did not have bleeds in the baseline lead-in period.

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Beqvez. Approval is recommended for those who meet the **Criteria** and **Dosing** for the listed indication. Because of the specialized skills required for evaluation and diagnosis of patients treated with Beqvez as well as the monitoring required for adverse events and long-term efficacy, approval requires Beqvez to be prescribed by a physician who specializes in the condition being treated. All approvals are provided for one-time (per lifetime) as a single dose. If claims history is available, verification is required for certain criteria as noted by **[verification in claims history required]**. For the dosing criteria, verification of the appropriate weight-based dosing is required by a Medical Director as noted by **[verification required]**. In the criteria for Beqvez, as appropriate, an asterisk (*) is noted next to the specified gender. In this context, the specified gender is defined as follows: males are defined as individuals with the biological traits of a man, regardless of the individual's gender identity or gender expression. All reviews (approvals and denials) will be forwarded to the Medical Director for evaluation.

Some clients have elected Embarc Benefit Protection. For these clients, the Medical Director will coordinate with EviCore to ensure the Embarc Benefit Protection portion of the review has been completed. If the Embarc Benefit Protection portion of the review has not been completed, the Medical Director will route to Embarc@eviCore.com prior to completing the review.

Documentation: Documentation is required for use of Beqvez as noted in the criteria as **[documentation required]**. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Beqvez is recommended in those who meet the following criteria:

FDA-Approved Indication

1. **Hemophilia B.** Approve a one-time (per lifetime) single dose if the patient meets ALL of the following (A, B, C, D, E, F, G, H, I, J, K, L, M, N, O, and P):
 - A) Patient is male*; AND
 - B) Patient is ≥ 18 years of age; AND
 - C) Patient has not received a gene therapy for hemophilia B in the past **[verification in claims history required]**; AND

Note: If no claim for Beqvez or Hemgenix (etranacogene dezaparvovec-drlb intravenous infusion) is present (or if claims history is not available), the prescribing physician confirms that the patient has not previously received Beqvez or Hemgenix.

- D) Patient has moderately severe or severe hemophilia B as evidenced by a baseline (without Factor IX replacement therapy) Factor IX level $\leq 2\%$ of normal **[documentation required]**; AND
- E) Patient meets ONE of the following (i, ii, or iii):
 - i. According to the prescribing physician, the patient has a history of use of Factor IX therapy for ≥ 150 exposure days; OR
 - ii. Patient meets BOTH of the following (a and b):
 - a) Patient has a history of life-threatening hemorrhage; AND
 - b) On-demand use of Factor IX therapy was required for this life-threatening hemorrhage; OR
 - iii. Patient meets BOTH of the following (a and b):
 - a) Patient has a history of repeated, serious spontaneous bleeding episodes; AND
 - b) On-demand use of Factor IX therapy was required for these serious spontaneous bleeding episodes; AND
- F) Patient does not have neutralizing antibodies to adeno-associated virus serotype Rh74var (AAVRh74var) capsid by an approved test **[documentation required]**; AND
- G) Patient meets BOTH of the following (i and ii):
 - i. Factor IX inhibitor titer testing has been performed within the past 30 days **[documentation required]**; AND
 - ii. Patient is negative for Factor IX inhibitors **[documentation required]**; AND
- H) Patient meets BOTH of the following (i and ii):
 - i. Patient does not have an active infection with hepatitis B virus or hepatitis C virus **[documentation required]**; AND
 - ii. Patient is not currently receiving antiviral therapy for a prior hepatitis B virus or hepatitis C virus exposure **[documentation required]**; AND
- I) According to the prescribing physician, the patient does not have uncontrolled human immunodeficiency virus infection; AND
- J) Patient has undergone liver function testing within the past 30 days and meets ALL of the following (i, ii, iii, and iv):
 - i. Alanine aminotransferase level is \leq two times the upper limit of normal **[documentation required]**; AND
 - ii. Aspartate aminotransferase level is \leq two times the upper limit of normal **[documentation required]**; AND
 - iii. Total bilirubin level is ≤ 1.5 times the upper limit of normal **[documentation required]**; AND
 - iv. Alkaline phosphatase level is \leq two times the upper limit of normal **[documentation required]**; AND
- K) Patient does not have evidence of advanced liver impairment and/or advanced fibrosis; AND
- L) Within the past 30 days, the platelet count was $\geq 100 \times 10^9/L$ **[documentation required]**; AND
- M) Within the past 30 days, creatinine was ≤ 2.0 mg/dL **[documentation required]**; AND
- N) The medication is prescribed by a hemophilia specialist physician; AND
- O) Current patient body weight has been obtained within the past 30 days **[documentation required]**; AND
- P) If criteria A through O are met, approve one dose (vials in a kit) of Beqvez to provide for a one-time (per lifetime) single dose of 5×10^{11} vector genomes per kg of body weight by intravenous infusion **[verification required]**. Table 1 provides the number of vials per kit and the National Drug Codes (NDCs) for each kit.
Note: Dose based on adjusted body weight for those with a body mass index $> 30 \text{ kg/m}^2$ using the following calculation: $\text{Dose Weight (kg)} = 30 \text{ kg/m}^2 \times [\text{Height (m)}]^2$

* Refer to the Policy Statement.

Dosing. The recommended dose of Beqvez is a one-time (per lifetime) single dose of 5×10^{11} vector genomes per kg of body weight by intravenous infusion.

Note: Dose based on adjusted body weight for those with a body mass index $> 30 \text{ kg/m}^2$ using the following calculation: $\text{Dose Weight (kg)} = 30 \text{ kg/m}^2 \times [\text{Height (m)}]^2$

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Beqvez is not recommended in the following situations:

- 1. Prior Receipt of Gene Therapy.** Prior receipt of gene therapy was a reason for patient exclusion in the pivotal study.
- 2. Patient with a History of Factor IX Inhibitors.** A history of Factor IX inhibitors was a reason for patient exclusion in the pivotal trial.
- 3.** Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

Table 1. Beqvez Multi-Vial Kits.¹

Patient Dose Weight	Total Number of Vials per Kit	NDC Number
$\leq 75 \text{ kg}$	4	0069-2004-04
$> 75 \text{ to } \leq 95 \text{ kg}$	5	0069-2005-05
$> 95 \text{ to } \leq 115 \text{ kg}$	6	0069-2006-06
$> 115 \text{ to } \leq 135 \text{ kg}$	7	0069-2007-07

NDC – National Drug Code.

REFERENCES

- Beqvez™ intravenous infusion [prescribing information]. New York, NY: Pfizer; April 2024.
- Cuker A, Kavakli K, Frenzel L, et al, for the BENEGENE-2 trial investigators. Gene therapy with fidanacogene elaparovvec in adults with hemophilia B. *N Engl J Med.* 2024;391(12):1108-1118.
- National Bleeding Disorders Foundation. Hemophilia B. An overview of symptoms, genetics, and treatments to help you understand hemophilia B. Available at: <https://www.hemophilia.org/bleeding-disorders-a-z/types/hemophilia-b>. Accessed on February 28, 2025.
- Sidonio RF, Malec L. Hemophilia (Factor IX deficiency). *Hematol Oncol Clin N Am.* 2021;35:1143-1155.
- Mancuso ME, Mahlangu JN, Pipe SW. The changing treatment landscape in haemophilia: from standard half-life clotting factor concentrates to gene editing. *Lancet.* 2021;397:630-640.
- Croteau SE. Hemophilia A/B. *Hematol Oncol Clin N Am.* 2022;36:797-812.
- Hympavzi™ subcutaneous injection [prescribing information]. New York, NY: Pfizer; October 2024.

HISTORY

Type of Revision	Summary of Changes	Review Date
New Policy	--	05/15/2024
UCare P&T Review	Policy reviewed and approved by UCare P&T committee. Annual review process	09/16/2024
Early Annual Revision	Hemophilia B: The requirement that the patient does not have a history of Factor IX inhibitors (with documentation required) was removed from this section. The requirement that prophylactic therapy with Factor IX will not be given after Beqvez administration once adequate Factor IX levels have been achieved was removed, along with the related Note. The Note that provides examples of advanced liver impairment and/or advanced fibrosis was removed. However, the criterion that the patient does not have evidence of advanced liver impairment and/or advanced fibrosis remains. Conditions Not Recommended for Approval: The condition of “Patient with a History of Factor IX Inhibitors” was added to this section. Previously, this was in a criterion related to the diagnosis of hemophilia B.	02/26/2025
UCare P&T Review	Policy reviewed and approved by UCare P&T committee. Annual review process	09/15/2025