

Utilization Review Policy 268

POLICY: Ophthalmology – Vascular Endothelial Growth Factor Inhibitors – Lucentis

- Lucentis® (ranibizumab intravitreal injection Genentech)
- Cimerli™ (ranibizumab-eqrn intravitreal injection Coherus)
- Byooviz[™] (ranibizumab-nuna intravitreal injection Biogen)

EFFECTIVE DATE: 1/1/2022

LAST REVISION DATE: 12/30/2024

COVERAGE CRITERIA FOR: All Aspirus Medicare Plans

OVERVIEW

Ranibizumab is a vascular endothelial growth factor (VEGF) inhibitor. There are two interchangeable biosimilars to Lucentis: Byooviz and Cimerli.

Lucentis and Cimerli are indicated for the following uses: 1,3

- Diabetic macular edema.
- Diabetic retinopathy.
- Macular edema following retinal vein occlusion.
- Myopic choroidal neovascularization.
- Neovascular (wet) age-related macular degeneration.

Byooviz is indicated for the following uses:²

- Macular edema following retinal vein occlusion.
- Myopic choroidal neovascularization.
- Neovascular (wet) age-related macular degeneration.

The recommended dosing for each of the indication is as follows: 1-3

- **Diabetic macular edema, diabetic retinopathy:** 0.3 mg administered by intravitreal injection once every month (approximately 28 days) [Cimerli and Lucentis)
- Macular edema following retinal vein occlusion, neovascular (wet) age-related macular degeneration: 0.5 mg administered by intravitreal injection once every month (approximately 28 days).
- **Myopic choroidal neovascularization:** 0.5 mg administered by intravitreal injection once every month (approximately 28 days) for up to 3 months; patients may be retreated if needed.

Other Uses with Supportive Evidence

Overproduction of VEGF may lead to other eye conditions, including neovascular glaucoma, retinopathy of prematurity, and other retinal and choroidal neovascular conditions affecting the eye.^{4,5} The VEGF inhibitors have the potential to be used off-label to reduce or slow visual impairment or vision loss associated with other eye conditions related to increased VEGF

production.^{4,6,7} The use of VEGF inhibitors have been shown to stop the angiogenic process, maintain visual acuity, and improve vision in patients with certain neovascular ophthalmic conditions. Therefore, research is rapidly evolving on the use of VEGF inhibitors in other neovascular ophthalmic conditions that threaten vision.

POLICY STATEMENT

Prior authorization is recommended for medical benefit coverage of ranibizumab products. Approval is recommended for those who meet the Criteria and Dosing for the listed indication(s). All approvals are provided for the duration noted below.

This policy incorporates Medicare coverage guidance as set forth in National Coverage Determinations (NCDs) and Local Coverage Determinations (LCDs), as well as in companion policy articles and other guidance applicable to the relevant service areas. These documents are cited in the References section of this policy. In some cases, this guidance includes specific lists of HCPCS and ICD-10 codes to help inform the coverage determination process. The Articles that include specific lists for billing and coding purposes will be included in the Reference section of this policy. However, to the extent that this policy cites such lists of HCPCS and ICD-10 codes, they should be used for reference purposes only. The presence of a specific HCPCS or ICD-10 code in a chart or companion article to an LCD is not by itself sufficient to approve coverage. Similarly, the absence of such a code does <u>not</u> necessarily mean that the applicable condition or diagnosis is excluded from coverage.

<u>Note</u>: Conditions for coverage outlined in this Medicare Advantage Medical Policy may be less restrictive than those found in applicable National Coverage Determinations, Local Coverage Determinations and/or Local Coverage Articles. Examples of situations where this clinical policy may be less restrictive include, but are not limited to, coverage of additional indications supported by CMS-approved compendia and the exclusion from this policy of additional coverage criteria requirements outlined in applicable National Coverage Determinations, Local Coverage Determinations and/or Local Coverage Articles.

Indications with a ^ below are also covered (and, if applicable, further detailed/referenced) in the corresponding Commercial Care Continuum (CC) Policy. Note: Additional criteria requirements for coverage of the same indication as outlined in the Commercial CC Policy and this Medicare Advantage CC Policy may NOT be the same.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Lucentis, Cimerli or Byooviz is recommended for requests meeting both the step therapy requirements and indication requirements:

Step Therapy Requirements (New Starts Only)

Criteria. The patient must meet the following criteria (A, B, or C):

- **A)** For patients new to Lucentis, Cimerli or Byooviz therapy only, must have a trial of repackaged Avastin prior to approval of Lucentis, Cimerli or Byooviz New starts to therapy defined as no use of Lucentis, Cimerli or Byooviz within the past 365 days for Medicare patients and includes use in either eye.
- **B)** Patient has diabetic retinopathy (without diabetic macular edema).

C) Patient has a contraindication or other clinical reason why repackaged Avastin cannot be tried before Lucentis, Cimerli or Byooviz.

Note: Step therapy only required for indications compendia supported for both Lucentis, Cimerli or Byooviz and Avastin.

FDA-Approved Indications

1. Neovascular (Wet) Age-Related Macular Degeneration. ^

Criteria. Approve for 1 year.

Dosing. Approve if the requested dosing meets the following (A <u>and</u> B):

- A) The dose is 0.5 mg administered by intravitreal injection for each eye being treated; AND
- **B)** The dosing interval is not more frequent than once every 28 days for each eye being treated.

2. Macular Edema Following Retinal Vein Occlusion. ^

Criteria. Approve for 1 year.

Dosing. Approve if the dose meets both criteria (A and B):

- A) The dose is 0.5 mg administered by intravitreal injection for each eye being treated; AND
- **B)** The dosing interval is not more frequent than once every 28 days for each eye being treated.

3. Diabetic Macular Edema. ^

Criteria. Approve for 1 year.

Dosing. Approve if the dose meets both criteria (A and B):

- A) The dose is 0.3 mg administered by intravitreal injection for each eye being treated; AND
- **B)** The dosing interval is not more frequent than once every 28 days for each eye being treated.

4. Diabetic Retinopathy. ^

Criteria. Approve for 1 year.

Dosing. Approve if the dose meets both criteria (A <u>and</u> B):

- A) The dose is 0.3 mg administered by intravitreal injection for each eye being treated; AND
- B) The dosing interval is not more frequent than once every 28 days for each eye being treated.

5. Myopic Choroidal Neovascularization. ^

Criteria. Approve for 1 year.

Dosing. Approve if the dose meets both criteria (A and B):

- A) The dose is 0.5 mg administered by intravitreal injection for each eye being treated; AND
- **B)** The dosing interval is not more frequent than once every 28 days for each eye being treated.

Other Uses with Supportive Evidence

6. Other Neovascular Diseases of the Eye. ^

<u>Note</u>: Examples of other neovascular diseases of the eye include neovascular glaucoma, retinopathy of prematurity, sickle cell neovascularization, choroidal neovascular conditions.

Criteria. Approve for 1 year.

Dosing. Approve if the dose meets both criteria (A <u>and</u> B):

- A) The dose is 0.5 mg administered by intravitreal injection for each eye being treated; AND
- **B)** The dosing interval is not more frequent than once every 28 days for each eye being treated.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of ranibizumab products is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

- 1. Lucentis® intravitreal injection [prescribing information]. South San Francisco, CA: Genentech; February 2024.
- 2. Byooviz[™] intravitreal injection [prescribing information]. Cambridge, MA: Biogen; October 2023.
- 3. Cimerli™ intravitreal injection [prescribing information]. Redwood City, CA: Coherus; May 2024.
- 4. Barakat MR, Kaiser PK. VEGF inhibitors for the treatment of neovascular age-related macular degeneration. *Expert Opin Investiq Drugs*. 2009;18(5):637-646.
- 5. Tolentino M. Systemic and ocular safety of intravitreal anti-VEGF therapies for ocular neovascular disease. *Surv Ophthalmol.* 2011;56(2):95-113.
- 6. Kinnunen K, Ylä-Herttuala S. Vascular endothelial growth factors in retinal and choroidal neovascular diseases. *Ann Med*. 2012;44(1):1-17.
- 7. Horsley MB, Kahook MY. Anti-VEGF therapy for glaucoma. Curr Opin Ophthalmol. 2010;21(2):112-117.
- 8. Centers for Medicare and Medicaid Services, National Government Services, Inc, Local Coverage Determination (LCD): Drugs and Biologicals, Coverage of, for Label and Off-Label Uses (L33394) [original date 10/01/2015; revision effective date 8/1/24]. Accessed on December 30, 2024.
- 9. Centers for Medicare and Medicaid Services, National Government Services, Inc, Local Coverage Article (LCA): Billing and Coding: Ranibizumab, Aflibercept and Brolucizumab-dbll (A52451) [original date 10/01/2015; revision effective date 9/1/24]. Accessed on December 30, 2024.

HISTORY

Type of Revision	Summary of Changes*	Date
Policy created	New Medicare Advantage Medical Care Continuum Policy	07/11/2018
Select revision	Added Macugen to policy	11/05/2018
Select revision	Reviewed and revised original policy created 07/11/2018 in	5/22/2019
	accordance with Local Coverage Article A52451 and Ophthalmology	

	Vascular Endothelial Growth Factor (VEGF) Inhibitor Injectables Care Continuous Hillington Review Religy	
Select revision	Care Continuum Utilization Review Policy. Reviewed and revised original policy created 07/11/2018 in	11/06/2019
Select levision	accordance with Local Coverage Determination L33394 and	11/00/2019
	Ophthalmology – Vascular Endothelial Growth Factor Inhibitors -	
	Lucentis Care Continuum Utilization Review Policy.	
Select revision	Non-clinical update to policy to add the statement "This policy	1/30/2020
	incorporates Medicare coverage guidance as set forth in National	
	Coverage Determinations (NCDs) and Local Coverage	
	Determinations (LCDs), as well as in companion policy articles and	
	other guidance applicable to the relevant service areas. These	
	documents are cited in the References section of this policy. In	
	some cases, this guidance includes specific lists of HCPCS and ICD-	
	10 codes to help inform the coverage determination process. The	
	Articles that include specific lists for billing and coding purposes will be included in the Reference section of this policy. However, to	
	the extent that this policy cites such lists of HCPCS and ICD-10	
	codes, they should be used for reference purposes only. The	
	presence of a specific HCPCS or ICD-10 code in a chart or	
	companion article to an LCD is not by itself sufficient to approve	
	coverage. Similarly, the absence of such a code does <u>not</u>	
	necessarily mean that the applicable condition or diagnosis is	
	excluded from coverage."	
Policy revision	*Added the following to the Policy Statement " <u>Note</u> : Conditions for	08/07/2020
	coverage outlined in this Medicare Advantage Medical Policy may	
	be less restrictive than those found in applicable National Coverage	
	Determinations, Local Coverage Determinations and/or Local	
	Coverage Articles. Examples of situations where this clinical policy	
	may be less restrictive include, but are not limited to, coverage of additional indications supported by CMS-approved compendia and	
	the exclusion from this policy of additional coverage criteria	
	requirements outlined in applicable National Coverage	
	Determinations, Local Coverage Determinations and/or Local	
	Coverage Articles."	
	*Updated references	
Selected	Product: Byooviz was added to the same conditions for approval	06/08/2022
Revision	as for Lucentis.	
Selected	Product: Cimerli was added to the same conditions for approval as	08/10/2022
Revision	for the other ranibizumab products. Clarified that continuation of therapy is acceptable if the requested	10/7/2022
Aspirus Revision	product has been used in either eye.	10/1/2022
Policy revision	For all indications/uses, the dosing interval was changed from "not	12/11/2023
Total Tevision	more frequent than once every 25 days for each eye being treated"	12,11,2323
	to "not more frequent than once every 28 days for each eye being	
	treated"; the 28 days aligns with the prescribing information.	
Policy review	No criteria changes, review based on LCD/LCA update.	09/11/2024
Aspirus P&T	Policy reviewed and approved by Aspirus P&T committee. Annual	09/16/2024
Review	review process	
Policy review	No criteria change, review based on commercial policy annual review	12/30/2024
Aspirus P&T Review	Policy reviewed and approved by Aspirus P&T committee. Annual review process	09/15/2025
	·	· · · · · · · · · · · · · · · · · · ·