

POLICY: Hemophilia – Gene Therapy – Roctavian Utilization Management Medical Policy

- Roctavian® (valoctocogene roxaparvovec-rvox intravenous infusion – BioMarin)

EFFECTIVE DATE: 11/15/2023

LAST REVISION DATE: 09/24/2025

COVERAGE CRITERIA FOR: All Aspirus Medicare Plans

OVERVIEW

Roctavian, an adeno-associated virus vector-based gene therapy, is indicated for the treatment of severe hemophilia A (congenital Factor VIII deficiency with Factor VIII activity < 1 IU/dL) in adults without pre-existing antibodies to adeno-associated virus serotype 5 detected by an FDA-approved test.¹

Disease Overview

Hemophilia A is an X-linked bleeding disorder primarily impacting males caused by a deficiency in Factor VIII.^{2,3} The incidence of hemophilia A in males is 1:5,000 livebirths.² The condition is characterized by prolonged bleeding after injuries, as well as spontaneous bleeds. Bleeding can occur in many different body areas as well (e.g., muscles, central nervous system, joints). The bleeding manifestations can lead to substantial morbidity such as hemophilic arthropathy.^{2,3} Disease severity is usually defined by the plasma levels or activity of Factor VIII classified as follows: severe (< 1%), moderate (1% to 5%), and mild (> 5% to < 40%); phenotypic expression may vary.³ The management of hemophilia involves use of agents to treat bleeding episodes, therapies given prophylactically to prevent bleeding, and gene therapy.^{2,3}

Clinical Efficacy

The efficacy of Roctavian was evaluated in one open-label, single-group, multinational Phase III trial (GENEr8-1) involving 134 adult males (\geq 18 years of age) with severe hemophilia A (Factor VIII activity level \leq 1 IU/dL).^{1,4,5} Patients involved in the trial did not have Factor VIII inhibitors (or a history of such inhibitors) and were receiving regular prophylaxis with Factor VIII products.⁴ Use of prophylactic Factor VIII therapy was not permitted during the trial, but could be used up to 4 weeks post Roctavian administration to allow the agent to have an effect.⁴ Other notable exclusion criteria were active infection, chronic or active hepatitis B or C, immunosuppressive disorder (including human immunodeficiency virus [HIV]), Stage 3 or 4 liver fibrosis, cirrhosis, liver function test abnormalities, serum creatinine \geq 1.4 mg/dL, and active malignancy.¹ Patients had to be treated or exposed to Factor VIII concentrates previously for a minimum of 150 exposure days.⁴ Among 132 patients (who were HIV-negative), the mean Factor VIII activity level at Weeks 49 through 52 had increased by 41.9 IU/dL.⁴ Among the 112 patients enrolled from the noninterventional study, the mean annualized rates of Factor VIII concentrate use and treated bleeding after Week 4 had decreased after Roctavian administration by 98.6% and 83.8%, respectively (P < 0.001 for both comparisons).^{4,5} At a median of 3 years post Roctavian dosing, the mean annualized bleeding rate in the rollover population (n = 112) in the efficacy

evaluation period was 2.6 bleeds/year compared to a mean baseline of 5.4 bleeds/year (while using Factor VIII therapies).¹ Long-term data (up to 4 years) reports favorable results with Roctavian.⁵⁻⁷

POLICY STATEMENT

Prior Authorization is recommended for medical benefit coverage of Roctavian. Approval is recommended for those who meet the **Criteria** and **Dosing** for the listed indication. Because of the specialized skills required for evaluation and diagnosis of patients treated with Roctavian as well as the monitoring required for adverse events and long-term efficacy, approval requires Roctavian to be prescribed by a physician who specializes in the condition being treated. All approvals are provided for one-time (per lifetime) as a single dose. If claims history is available, verification is required for certain criteria as noted by **[verification in claims history required]**. For the dosing criteria, verification of the appropriate weight-based dosing is required by a Medical Director as noted by **[verification required]**. In the criteria for Roctavian, as appropriate, an asterisk (*) is noted next to the specified gender. In this context, the specified gender is defined as follows: males are defined as individuals with the biological traits of a man, regardless of the individual's gender identity or gender expression. All reviews (approvals and denials) will be forwarded to the Medical Director for evaluation.

Some clients have elected Embarc Benefit Protection. For these clients, the Medical Director will coordinate with EviCore to ensure the Embarc Benefit Protection portion of the review has been completed. If the Embarc Benefit Protection portion of the review has not been completed, the Medical Director will route to Embarc@EviCore.com prior to completing the review.

Documentation: Documentation is required for use of Roctavian as noted in the criteria as **[documentation required]**. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Roctavian is recommended in those who meet the following criteria:

FDA-Approved Indication

-
- 1. Hemophilia A.** Approve a one-time (per lifetime) single dose if the patient meets ALL of the following (A, B, C, D, E, F, G, H, I, J, K, L, M, N, O, P, Q, R, and S):
 - A)** Patient is male* AND
 - B)** Patient is ≥ 18 years of age; AND
 - C)** Patient has not received Roctavian in the past **[verification in claims history required]**; AND **Note:** If no claim for Roctavian is present (or if claims history is not available), the prescribing physician confirms that the patient has not previously received Roctavian.
 - D)** Patient has severe hemophilia A as evidence by a baseline (without Factor VIII replacement therapy) Factor VIII level of < 1 IU/dL **[documentation required]**; AND
-

- E) Patient does not have detectable pre-existing antibodies to adeno-associated virus 5 (AAV5) by an FDA-approved test **[documentation required]**; AND
- F) According to the prescribing physician, the patient has a history of use of Factor VIII therapy for at least 150 exposure days; AND
- G) Patient meets ALL of the following (i, ii, and iii):
 - i. Factor VIII inhibitor titer testing has been performed within the past 30 days **[documentation required]**; AND
 - ii. Patient does not currently have an inhibitor to Factor VIII **[documentation required]**; AND
 - iii. Patient does not have a history of Factor VIII inhibitors **[documentation required]**; AND
- H) Prophylactic therapy with Factor VIII will not be given after Roctavian administration once adequate Factor VIII levels have been achieved; AND
Note: Use of episodic Factor VIII therapy is acceptable for the treatment of bleeds and for surgery/procedures if needed as determined by the hemophilia specialist physician.
- I) Patient does not have a known hypersensitivity to mannitol; AND
- J) Patient does not have chronic or active hepatitis B **[documentation required]**; AND
- K) Patient does not have active hepatitis C **[documentation required]**; AND
- L) Patient is not human immunodeficiency virus positive **[documentation required]**; AND
- M) Patient does not have evidence of significant hepatic fibrosis or cirrhosis; AND
- N) Patient meets ONE of the following (i or ii):
 - i. Patient has undergone liver function testing within the past 30 days and meets ALL of the following (a, b, c, d, e, and f):
 - a) Alanine aminotransferase levels are ≤ 1.25 times the upper limit of normal **[documentation required]**; AND
 - b) Aspartate aminotransferase levels are ≤ 1.25 times the upper limit of normal **[documentation required]**; AND
 - c) Total bilirubin levels are ≤ 1.25 times the upper limit of normal **[documentation required]**; AND
 - d) Alkaline phosphatase levels are ≤ 1.25 times the upper limit of normal **[documentation required]**; AND
 - e) Gamma-glutamyl transferase levels are ≤ 1.25 times the upper limit of normal **[documentation required]**; AND
 - f) The International Normalized Ratio is < 1.4 **[documentation required]**; OR
 - ii. If the patient had one or more of the laboratory values listed in *Criteria a-f* above that was not at the value specified in *Criteria a-f* above, then a hepatologist has evaluated the patient and has determined that use of Roctavian is clinically appropriate **[documentation required]**; AND
- O) Within the past 30 days, the platelet count was $\geq 100 \times 10^9/L$ **[documentation required]**; AND
- P) Within the past 30 days, the creatinine level was < 1.4 mg/dL **[documentation required]**; AND
- Q) Medication is prescribed by a hemophilia specialist physician; AND
- R) Current patient body weight has been obtained within the past 30 days **[documentation required]**; AND
- S) If criteria A through R are met, approve one dose of Roctavian to provide for a one-time (per lifetime) single dose of 6×10^{13} vector genomes per kg by intravenous infusion **[verification required]**.

Note: Roctavian is supplied in a carton (NDC 68135-927-48) that contains one single dose vial (NDC 68135-927-01) with an extractable volume of not less than 8 mL, containing 16×10^{13} vector genomes.

* Refer to the Policy Statement.

Dosing. The recommended dose of Roctavian is a one-time (per lifetime) single dose of 6×10^{13} vector genomes per kg based on body weight in kg by intravenous infusion.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Roctavian is not recommended in the following situations:

- 1. Prior Receipt of Gene Therapy.** Prior receipt of gene therapy was a reason for patient exclusion in the pivotal study.
- Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

- Roctavian® intravenous infusion [prescribing information]. Novato, CA: BioMarin; June 2023.
- Chowdary P, Carcao M, Kenet G, Pipe W. Haemophilia. *Lancet*. 2025;405:736-750.
- Konkle BA, Nakaya Fletcher S. Hemophilia A. 2000 Sep 21 [updated 2025 Aug 7]. In: Adam MP, Feldman J, Mirzaa GM, Pagon RA, Wallace SE, Amemiya A, editors. *GeneReviews*® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2025. PMID: 20301578.
- Ozelo MC, Mahlangu J, Pasi KJ, et al, for the GENER8-1 trial group. Valoctocogene roxaparovec gene therapy for hemophilia A. *N Engl J Med*. 2022;386(11):1013-1025.
- Mahlangu J, Kaczmarek R, Von Drygalski A, et al, for the GENER8-1 trial group. Two-year outcomes of valoctocogene roxaparovec therapy for hemophilia A. *N Engl J Med*. 2023;388(8):694-705.
- Madan B, Ozelo MC, Paheja P, et al. Three-year outcomes of valoctocogene roxaparovec gene therapy for hemophilia A. *J Thromb Haemost*. 2024;22:1880-1893.
- Leavitt AD, Mahlangu J, Raheja P, et al. Efficacy, safety, and quality of life 4 years after valoctocogene roxaparovec gene transfer for severe hemophilia A in the phase 3 GENER8-1 trial. *Res Pract Thromb Haemost*. 2024;8:e102615.

HISTORY

Type of Revision	Summary of Changes	Review Date
New Policy	--	08/16/2023
Annual Revision	The Policy Statement was clarified to add that all approvals are provided for one-time (per lifetime) as a single dose. The phrase “if claims history is available” was added regarding that verification in claims history is required for certain criteria. Regarding Documentation, “medical test results” and “prescription receipts” were added; laboratory “tests” was changed to “results”. In addition, the following changes were made: Hemophilia A: <ul style="list-style-type: none"> For approval, the word “single” was added before the word “dose” for clarification. Regarding the Note in the criteria which addresses that the patient has not received Roctavian in the past (with verification in claims history required), a phrase was added to include situations in which claims history is not available. 	09/11/2024

	<ul style="list-style-type: none"> • The phrase “according to the prescribing physician” was added to the requirement that the patient has a history of use of Factor VIII therapy for at least 150 exposure days. • The phrase “within 30 days before the intended receipt of Roctavian” was replaced with “within the past 30 days” regarding the requirement that Factor VIII inhibitor titer testing has been performed. • The requirement was removed that the patient does not have an active acute or uncontrolled chronic infection. • The phrase “liver health assessment” was replaced with “liver function testing”. Also, the phrase “within 30 days before the intended receipt of Roctavian” was replaced with “within the past 30 days” regarding the liver function testing. • The phrase, “within 30 days before the intended receipt of Roctavian” was replaced with “within the past 30 days” regarding the requirement that the platelet count was $\geq 100 \times 10^9/L$. • The phrase “within 30 days before the intended receipt of Roctavian” was replaced with “within the past 30 days” regarding the requirement that the creatinine level was $< 1.4 \text{ mg/dL}$. • The requirement (along with the related Note) was removed that the patient has not used a systemic immunosuppressive agent within 30 days before intended receipt of Roctavian. • The requirement was removed that the patient does not have any disease or condition that would interfere with the compliance requirements that involve use of systemic corticosteroid therapy or systemic alternative immunosuppressive medications. • The requirement was removed that the patient does not have an immunosuppressive disorder. • The requirement was removed that the patient does not have any additional bleeding disorder, besides hemophilia A. • The requirement was removed that the patient does not have a history of thrombosis or thrombophilia. • The requirement (along with the related Note) was removed that the patient does not have a current active malignancy. • The requirement was removed that the patient does not have a history of hepatic malignancy. • The requirement was removed that the patient has not received a live vaccine within 30 days before intended receipt of Roctavian. • The requirement was removed that the hemophilia specialist physician has discussed with the patient that for a period of up to 6 months after administration of Roctavian that precautions should be taken that a male of reproductive potential (and his female partner) prevent or postpone pregnancy by utilizing an effective form of contraception and that a male should not donate semen. • The phrase “within 30 days before the intended receipt of Roctavian” was replaced with “within the past 30 days” regarding the requirement that current patient body weight has been obtained. <p>Conditions Not Recommended for Approval: The condition of “Prior Receipt of Gene Therapy” was added.</p>	
Aspirus P&T Review	Policy reviewed and approved by Aspirus P&T committee. Annual review process	09/16/2024
Aspirus P&T Review	Policy reviewed and approved by Aspirus P&T committee. Annual review process	09/15/2025
Annual Revision	No criteria changes.	09/24/2025